

## **WARNING**

**Harmful if swallowed**

**Harmful in contact with skin**

**Keep out of reach of children**

## **RAPPEL®**

### **Seed Applied Slug Repellent**

**Contains: Metaldehyde, Fulvic acid, Humic acid and Saponin**

#### ***Seed Applied Slug Repellent***

**A seed treatment directly applied to forage brassica, pasture, cereal and maize seed for the purpose of repelling slugs during plant establishment.**

#### **Precautions:**

Users must take all reasonable steps to ensure that RAPPEL® treated seed does not cause any significant adverse effects to the environment beyond the application area.

Treated seed is not to be used for human or animal consumption.

- Withholding Period:** Stock must not graze for 42 days following sowing and for grass seed oversown into pasture (stitching in) within 21 days from sowing.
- Storage:** Treated seed should be stored out of the reach of animals and children.
- Sowing:** Treated seed may be harmful to seed eating birds. It should be properly sown and covered by soil except when used in combination with a suitable bird repellent.
- Handling:** Gloves and a dust mask should be worn when handling seed. Exposed skin should be flushed in running cold water after handling seed.
- Disposal:** Any excess seed should be removed from the environment and rebagged into the original bags.

Avoid contamination of any water supply or water body with empty packaging or excess product.

Do not reuse packaging. Dispose of packaging responsibly.

**Bee Safety:** Do not broadcast or drill RAPPEL® treated seed into paddocks where bees are foraging.

**Use by Date:** Use within 1 year from date of purchase.

### **Directions for Use:**

This product is not for sale as a seed treatment formulation but as a custom treatment on seed from accredited applicators.

Sow RAPPEL® treated forage brassica, pasture, cereal and maize seed at normal commercial rates where there is a risk of slug damage during establishment.

Ensure RAPPEL® treated seed is properly sown and covered by soil except when used in combination with a suitable bird repellent.

Continue to monitor for slug damage during establishment as high numbers of slugs may overwhelm the repellent effect of the treatment and further actions may be required.

### **Date of Manufacture:**

### **Batch Number:**

**Manufactured by:** Hodder & Taylors Ltd  
69 Kawakawa Road  
Feilding 4740

Ph. 06 3235091  
Email: [info@hnt.co.nz](mailto:info@hnt.co.nz)